

Fibre supplements act by increasing the amount of dietary fibre, drawing some fluid into the bowel and bulking up stools. Various commercial preparations are available including fibre husks, tablets and capsules and dissolvable powders. Excess fibre intake may cause abdominal discomfort, bloating and flatulence. Fibre supplements alone however tend to be insufficient to treat moderate to severe constipation and can

Stool softeners act by drawing fluid into the intestine to soften the stools. Many preparations used to clean the bowel out before colonoscopy also belong to this group of laxatives. There are several different types of stool softeners including unabsorbable sugars (e.g. sorbitol, lactulose), magnesium, phosphate or citrate salts, polyethylene glycol and the newer macrogol preparations. These preparations are available as syrup, powders or enemas. Side effects of these include bloating, flatulence, abdominal cramps, explosive diarrhoea

### Bowel Stimulants

Stimulant laxatives are those that help the bowel to contract and pass stools along. These include senna, cascara and bisacodyl; and are available as granules, tablets, fruit pastes or suppositories. Castor oil which was popular many years ago is both a lubricant and stimulant of the bowel and is now rarely used. Liquorice, caffeine and nicotine also have some stimulant effects on the bowel, but on their own are usually insufficient to treat constipation. "Natural" laxatives including juices and fruit

Dr. Rela Institute & Medical Centre is a multi-specialty quaternary care hospital located in Chennai, India.

The Institute is within the campus of Sree Balaji Medical College and Hospital, which is spread across 36 acres. It has 14 operating theatres with 450 beds, inclusive of 150 critical care beds.

The Institute is conveniently located 10 minutes from the Domestic and International Airport.

The hospital is designed to provide highly specialized care in various departments with a focus on multi-organ transplantation. Prof. Mohamed Rela, a world renowned surgeon in the field of Liver surgery and transplantation is the Chairman and Managing Director of the Institute.

In addition to quaternary & quality care, is also committed to provide day to day primary and secondary care to the local population, with facilities of international standards.

The Institute would provide comprehensive support to international patients travelling for medical treatment

24 Hr Emergency ☎ 044-6666 7788



**DR. RELA INSTITUTE & MEDICAL CENTRE**

No. 7, CLC Works Road, Chromepet,  
Chennai - 600 044, Tamil Nadu, INDIA.

Tel : +91 44 6666 7777

Email: helpdesk@relainstitute.com

[www.relainstitute.com](http://www.relainstitute.com)



**DR. RELA INSTITUTE & MEDICAL CENTRE**  
An International Medical Facility



Department of  
Gastroenterology

## Constipation

# Constipation

## 1 What is constipation?

Constipation is a common medical complaint. One in 10 children will seek medical attention for this problem, which accounts for 3-5% of all visits to a paediatrician. It is also a common complaint in adults, particularly women. Up to one in five adults report that they have constipation. It increases with advancing age, particularly in those over 65 years of age. People say they are constipated when they can't empty their bowels as often or as easily as they would like. There is however a wide variation as to what is considered normal which can range from

2 Constipation can be functional (i.e. no underlying disease) or may result from other causes (secondary constipation). Symptoms such as a recent change in bowel habits, weight loss, persistent or profuse bleeding, or severe abdominal pain suggest that there may be an underlying cause for the constipation. Use of certain drugs and medications, hormonal and metabolic disorders, neurological disorders (e.g. Parkinson's disease, stroke) and pregnancy can also result in constipation. Many women also find they tend to be constipated just before their menstrual periods as hormonal changes can also affect their bowels. In infants and young children, cow's milk allergy has been recognised to cause severe constipation in a minority of children. Most patients, especially those who have had symptoms for years, have functional constipation and do not have any underlying serious disease. This condition often runs in families and almost a third of patients will have

## 3 What symptoms are associated with constipation?

The commonest complaint is that of infrequent bowel motions, usually fewer than three times per week. Other complaints

o

o

o The feeling of incomplete evacuation or of a blockage

Some people with severe symptoms also have to use special manoeuvres (e.g. pushing on the perineum or using their fingers) to help them do a poo.

If stools are very hard, dry and large, it may cause a small split in the lining of the back passage (anal fissure) and result in pain and stinging in the area as well as some bright red bleeding. In those with very long standing constipation, particularly children and the elderly, the lower part of the bowel (rectum) may become stretched. This will often result in loss of awareness of the need to pass a motion and cause faecal

## 4 Does diet play a role?

Promoting a healthy diet with adequate fluid and fibre is helpful and may be all that is required in some patients to regulate and manage their bowels.

The recommended fibre intake:

Adults: 25 to 30g daily

Children: Child's age plus 5g / day (e.g. a 7yr old should have 7+5 = 12g/day)

In those unable to increase their fibre intake from dietary changes, commercial fibre supplements are available and can

## 5

The best times to try to pass motions are on waking in the morning or shortly after meals. This is because the bowel is most active then.

When sitting on the toilet it may be useful to make sure the knees are bent above the level of the hips and the feet are flat on the floor (Indian Style). A small foot-stool under the feet may be necessary to get the best position in Western style. Reading, doing puzzles and playing electronic games are generally not recommended while attempting to have bowel motions because it tends to significantly prolong toileting time and

## 6

Laxative use, even for long periods of time, is safe. If however you need to take laxatives regularly to pass motions, you should seek medical attention to make sure there is no underlying cause for the constipation. Those who continue to have problems even with regular laxative use may need to see a specialist gastroenterologist.

Many healthy children have problems with constipation,

## 7 Do you have a good diet?

Aim for:

Brown or wholemeal breads and high fibre cereals - 4 or more serves/Day

Dairy products - 3 serves

Fruit - 2 or 3 pieces

Vegetables - 5 serves

Meat, Cheese, Poultry, fish - 1 or 2 serves/day

## 8

There are many different types of laxative preparations available, ranging from tablets, powders, syrups, fruit pastes to local preparations inserted directly into the back passage (enema or suppository). There are three main types of

- o Bulking agents including fibre supplements
- o Stool softeners and lubricants, and
- o Bowel stimulants.